


10.3 Two-Way Tables and Probability

Essential Question How can you construct and interpret a two-way table?

EXPLORATION 1 Completing and Using a Two-Way Table

Work with a partner. A *two-way table* displays the same information as a Venn diagram. In a two-way table, one category is represented by the rows and the other category is represented by the columns.



The Venn diagram shows the results of a survey in which 80 students were asked whether they play a musical instrument and whether they speak a foreign language. Use the Venn diagram to complete the two-way table. Then use the two-way table to answer each question.

	Play an Instrument	Do Not Play an Instrument	Total
Speak a Foreign Language			
Do Not Speak a Foreign Language			
Total			

- How many students play an instrument?
- How many students speak a foreign language?
- How many students play an instrument and speak a foreign language?
- How many students do not play an instrument and do not speak a foreign language?
- How many students play an instrument and do not speak a foreign language?

EXPLORATION 2 Two-Way Tables and Probability

Work with a partner. In Exploration 1, one student is selected at random from the 80 students who took the survey. Find the probability that the student

- plays an instrument.
- speaks a foreign language.
- plays an instrument and speaks a foreign language.
- does not play an instrument and does not speak a foreign language.
- plays an instrument and does not speak a foreign language.

EXPLORATION 3 Conducting a Survey

Work with your class. Conduct a survey of the students in your class. Choose two categories that are different from those given in Explorations 1 and 2. Then summarize the results in both a Venn diagram and a two-way table. Discuss the results.

MODELING WITH MATHEMATICS

To be proficient in math, you need to identify important quantities in a practical situation and map their relationships using such tools as diagrams and two-way tables.

Communicate Your Answer

- How can you construct and interpret a two-way table?
- How can you use a two-way table to determine probabilities?

10.3 Lesson

Core Vocabulary

two-way table, p. 554
 joint frequency, p. 554
 marginal frequency, p. 554
 joint relative frequency, p. 555
 marginal relative frequency, p. 555
 conditional relative frequency, p. 555

Previous
 conditional probability

READING

A two-way table is also called a *contingency table*, or a *two-way frequency table*.

What You Will Learn

- ▶ Make two-way tables.
- ▶ Find relative and conditional relative frequencies.
- ▶ Use conditional relative frequencies to find conditional probabilities.

Making Two-Way Tables

A **two-way table** is a frequency table that displays data collected from one source that belong to two different categories. One category of data is represented by rows and the other is represented by columns. Suppose you randomly survey freshmen and sophomores about whether they are attending a school concert. A two-way table is one way to organize your results.

Each entry in the table is called a **joint frequency**. The sums of the rows and columns are called **marginal frequencies**, which you will find in Example 1.

		Attendance	
		Attending	Not Attending
Class	Freshman	25	44
	Sophomore	80	32

joint frequency

EXAMPLE 1 Making a Two-Way Table

In another survey similar to the one above, 106 juniors and 114 seniors respond. Of those, 42 juniors and 77 seniors plan on attending. Organize these results in a two-way table. Then find and interpret the marginal frequencies.

SOLUTION

- Step 1** Find the joint frequencies. Because 42 of the 106 juniors are attending, $106 - 42 = 64$ juniors are not attending. Because 77 of the 114 seniors are attending, $114 - 77 = 37$ seniors are not attending. Place each joint frequency in its corresponding cell.
- Step 2** Find the marginal frequencies. Create a new column and row for the sums. Then add the entries and interpret the results.

		Attendance		Total
		Attending	Not Attending	
Class	Junior	42	64	106
	Senior	77	37	114
Total		119	101	220

119 students are attending. 101 students are not attending.

106 juniors responded.
114 seniors responded.
220 students were surveyed.

- Step 3** Find the sums of the marginal frequencies. Notice the sums $106 + 114 = 220$ and $119 + 101 = 220$ are equal. Place this value at the bottom right.

Monitoring Progress Help in English and Spanish at BigIdeasMath.com

1. You randomly survey students about whether they are in favor of planting a community garden at school. Of 96 boys surveyed, 61 are in favor. Of 88 girls surveyed, 17 are against. Organize the results in a two-way table. Then find and interpret the marginal frequencies.

Finding Relative and Conditional Relative Frequencies

You can display values in a two-way table as frequency counts (as in Example 1) or as *relative frequencies*.

Core Concept

STUDY TIP

Two-way tables can display relative frequencies based on the total number of observations, the row totals, or the column totals.

Relative and Conditional Relative Frequencies

A **joint relative frequency** is the ratio of a frequency that is not in the total row or the total column to the total number of values or observations.

A **marginal relative frequency** is the sum of the joint relative frequencies in a row or a column.

A **conditional relative frequency** is the ratio of a joint relative frequency to the marginal relative frequency. You can find a conditional relative frequency using a row total or a column total of a two-way table.

EXAMPLE 2 Finding Joint and Marginal Relative Frequencies

Use the survey results in Example 1 to make a two-way table that shows the joint and marginal relative frequencies.

SOLUTION

To find the joint relative frequencies, divide each frequency by the total number of students in the survey. Then find the sum of each row and each column to find the marginal relative frequencies.

		Attendance		Total
		Attending	Not Attending	
Class	Junior	$\frac{42}{220} \approx 0.191$	$\frac{64}{220} \approx 0.291$	0.482
	Senior	$\frac{77}{220} = 0.35$	$\frac{37}{220} \approx 0.168$	0.518
Total		0.541	0.459	1

About 29.1% of the students in the survey are juniors and are *not* attending the concert.

About 51.8% of the students in the survey are seniors.

INTERPRETING MATHEMATICAL RESULTS

Relative frequencies can be interpreted as probabilities. The probability that a randomly selected student is a junior and is *not* attending the concert is 29.1%.

EXAMPLE 3 Finding Conditional Relative Frequencies

Use the survey results in Example 1 to make a two-way table that shows the conditional relative frequencies based on the row totals.

SOLUTION

Use the marginal relative frequency of each *row* to calculate the conditional relative frequencies.

		Attendance	
		Attending	Not Attending
Class	Junior	$\frac{0.191}{0.482} \approx 0.396$	$\frac{0.291}{0.482} \approx 0.604$
	Senior	$\frac{0.35}{0.518} \approx 0.676$	$\frac{0.168}{0.518} \approx 0.324$

Given that a student is a senior, the conditional relative frequency that he or she is *not* attending the concert is about 32.4%.

- Use the survey results in Monitoring Progress Question 1 to make a two-way table that shows the joint and marginal relative frequencies.
- Use the survey results in Example 1 to make a two-way table that shows the conditional relative frequencies based on the column totals. Interpret the conditional relative frequencies in the context of the problem.
- Use the survey results in Monitoring Progress Question 1 to make a two-way table that shows the conditional relative frequencies based on the row totals. Interpret the conditional relative frequencies in the context of the problem.

Finding Conditional Probabilities

You can use conditional relative frequencies to find conditional probabilities.

EXAMPLE 4 Finding Conditional Probabilities

A satellite TV provider surveys customers in three cities. The survey asks whether they would recommend the TV provider to a friend. The results, given as joint relative frequencies, are shown in the two-way table.

		Location		
		Glendale	Santa Monica	Long Beach
Response	Yes	0.29	0.27	0.32
	No	0.05	0.03	0.04

- What is the probability that a randomly selected customer who is located in Glendale will recommend the provider?
- What is the probability that a randomly selected customer who will not recommend the provider is located in Long Beach?
- Determine whether recommending the provider to a friend and living in Long Beach are independent events.

SOLUTION

INTERPRETING MATHEMATICAL RESULTS

The probability 0.853 is a conditional relative frequency based on a column total. The condition is that the customer lives in Glendale.

$$\text{a. } P(\text{yes} | \text{Glendale}) = \frac{P(\text{Glendale and yes})}{P(\text{Glendale})} = \frac{0.29}{0.29 + 0.05} \approx 0.853$$

- So, the probability that a customer who is located in Glendale will recommend the provider is about 85.3%.

$$\text{b. } P(\text{Long Beach} | \text{no}) = \frac{P(\text{no and Long Beach})}{P(\text{no})} = \frac{0.04}{0.05 + 0.03 + 0.04} \approx 0.333$$

- So, the probability that a customer who will not recommend the provider is located in Long Beach is about 33.3%.

- c. Use the formula $P(B) = P(B|A)$ and compare $P(\text{Long Beach})$ and $P(\text{Long Beach} | \text{yes})$.

$$P(\text{Long Beach}) = 0.32 + 0.04 = 0.36$$

$$P(\text{Long Beach} | \text{yes}) = \frac{P(\text{Yes and Long Beach})}{P(\text{yes})} = \frac{0.32}{0.29 + 0.27 + 0.32} \approx 0.36$$

- Because $P(\text{Long Beach}) \approx P(\text{Long Beach} | \text{yes})$, the two events are independent.

Monitoring Progress Help in English and Spanish at BigIdeasMath.com

- In Example 4, what is the probability that a randomly selected customer who is located in Santa Monica will not recommend the provider to a friend?
- In Example 4, determine whether recommending the provider to a friend and living in Santa Monica are independent events. Explain your reasoning.

EXAMPLE 5 Comparing Conditional Probabilities



A jogger wants to burn a certain number of calories during his workout. He maps out three possible jogging routes. Before each workout, he randomly selects a route, and then determines the number of calories he burns and whether he reaches his goal. The table shows his findings. Which route should he use?

	Reaches Goal	Does Not Reach Goal
Route A		
Route B		
Route C		

SOLUTION

Step 1 Use the findings to make a two-way table that shows the joint and marginal relative frequencies. There are a total of 50 observations in the table.

Step 2 Find the conditional probabilities by dividing each joint relative frequency in the “Reaches Goal” column by the marginal relative frequency in its corresponding row.

		Result		Total
		Reaches Goal	Does Not Reach Goal	
Route	A	0.22	0.12	0.34
	B	0.22	0.08	0.30
	C	0.24	0.12	0.36
Total		0.68	0.32	1

$$P(\text{reaches goal} | \text{Route A}) = \frac{P(\text{Route A and reaches goal})}{P(\text{Route A})} = \frac{0.22}{0.34} \approx 0.647$$

$$P(\text{reaches goal} | \text{Route B}) = \frac{P(\text{Route B and reaches goal})}{P(\text{Route B})} = \frac{0.22}{0.30} \approx 0.733$$

$$P(\text{reaches goal} | \text{Route C}) = \frac{P(\text{Route C and reaches goal})}{P(\text{Route C})} = \frac{0.24}{0.36} \approx 0.667$$

- Based on the sample, the probability that he reaches his goal is greatest when he uses Route B. So, he should use Route B.

Monitoring Progress Help in English and Spanish at BigIdeasMath.com

- A manager is assessing three employees in order to offer one of them a promotion. Over a period of time, the manager records whether the employees meet or exceed expectations on their assigned tasks. The table shows the manager’s results. Which employee should be offered the promotion? Explain.

	Exceed Expectations	Meet Expectations
Joy		
Elena		
Sam		

Vocabulary and Core Concept Check

- COMPLETE THE SENTENCE** A(n) _____ displays data collected from the same source that belongs to two different categories.
- WRITING** Compare the definitions of joint relative frequency, marginal relative frequency, and conditional relative frequency.

Monitoring Progress and Modeling with Mathematics

In Exercises 3 and 4, complete the two-way table.

3.

		Preparation		Total
		Studied	Did Not Study	
Grade	Pass		6	
	Fail			10
Total		38		50

4.

		Response		Total
		Yes	No	
Role	Student	56		
	Teacher		7	10
Total			49	

5. **MODELING WITH MATHEMATICS** You survey 171 males and 180 females at Grand Central Station in New York City. Of those, 132 males and 151 females wash their hands after using the public rest rooms. Organize these results in a two-way table. Then find and interpret the marginal frequencies. (See Example 1.)



6. **MODELING WITH MATHEMATICS** A survey asks 60 teachers and 48 parents whether school uniforms reduce distractions in school. Of those, 49 teachers and 18 parents say uniforms reduce distractions in school. Organize these results in a two-way table. Then find and interpret the marginal frequencies.

USING STRUCTURE In Exercises 7 and 8, use the two-way table to create a two-way table that shows the joint and marginal relative frequencies.

7.

		Dominant Hand		Total
		Left	Right	
Gender	Female	11	104	115
	Male	24	92	116
Total		35	196	231

8.

		Gender		Total
		Male	Female	
Experience	Expert	62	6	68
	Average	275	24	299
	Novice	40	3	43
Total		377	33	410

9. **MODELING WITH MATHEMATICS** Use the survey results from Exercise 5 to make a two-way table that shows the joint and marginal relative frequencies. (See Example 2.)
10. **MODELING WITH MATHEMATICS** In a survey, 49 people received a flu vaccine before the flu season and 63 people did not receive the vaccine. Of those who receive the flu vaccine, 16 people got the flu. Of those who did not receive the vaccine, 17 got the flu. Make a two-way table that shows the joint and marginal relative frequencies.



11. MODELING WITH MATHEMATICS A survey finds that 110 people ate breakfast and 30 people skipped breakfast. Of those who ate breakfast, 10 people felt tired. Of those who skipped breakfast, 10 people felt tired. Make a two-way table that shows the conditional relative frequencies based on the breakfast totals. (See Example 3.)

12. MODELING WITH MATHEMATICS Use the survey results from Exercise 10 to make a two-way table that shows the conditional relative frequencies based on the flu vaccine totals.

13. PROBLEM SOLVING Three different local hospitals in New York surveyed their patients. The survey asked whether the patient's physician communicated efficiently. The results, given as joint relative frequencies, are shown in the two-way table. (See Example 4.)

		Location		
		Glens Falls	Saratoga	Albany
Response	Yes	0.123	0.288	0.338
	No	0.042	0.077	0.131

- What is the probability that a randomly selected patient located in Saratoga was satisfied with the communication of the physician?
- What is the probability that a randomly selected patient who was not satisfied with the physician's communication is located in Glens Falls?
- Determine whether being satisfied with the communication of the physician and living in Saratoga are independent events.

14. PROBLEM SOLVING A researcher surveys a random sample of high school students in seven states. The survey asks whether students plan to stay in their home state after graduation. The results, given as joint relative frequencies, are shown in the two-way table.

		Location		
		Nebraska	North Carolina	Other States
Response	Yes	0.044	0.051	0.056
	No	0.400	0.193	0.256

- What is the probability that a randomly selected student who lives in Nebraska plans to stay in his or her home state after graduation?
- What is the probability that a randomly selected student who does not plan to stay in his or her home state after graduation lives in North Carolina?
- Determine whether planning to stay in their home state and living in Nebraska are independent events.

ERROR ANALYSIS In Exercises 15 and 16, describe and correct the error in finding the given conditional probability.

		City			Total
		Tokyo	London	Washington, D.C.	
Response	Yes	0.049	0.136	0.171	0.356
	No	0.341	0.112	0.191	0.644
Total		0.39	0.248	0.362	1

15. $P(\text{yes} | \text{Tokyo})$

X

$$P(\text{yes} | \text{Tokyo}) = \frac{P(\text{Tokyo and yes})}{P(\text{Tokyo})}$$

$$= \frac{0.049}{0.356} \approx 0.138$$

16. $P(\text{London} | \text{no})$

X

$$P(\text{London} | \text{no}) = \frac{P(\text{no and London})}{P(\text{London})}$$

$$= \frac{0.112}{0.248} \approx 0.452$$

17. PROBLEM SOLVING You want to find the quickest route to school. You map out three routes. Before school, you randomly select a route and record whether you are late or on time. The table shows your findings. Assuming you leave at the same time each morning, which route should you use? Explain. (See Example 5.)

	On Time	Late
Route A		
Route B		
Route C		

18. PROBLEM SOLVING A teacher is assessing three groups of students in order to offer one group a prize. Over a period of time, the teacher records whether the groups meet or exceed expectations on their assigned tasks. The table shows the teacher's results. Which group should be awarded the prize? Explain.

	Exceed Expectations	Meet Expectations
Group 1		
Group 2		
Group 3		

19. **OPEN-ENDED** Create and conduct a survey in your class. Organize the results in a two-way table. Then create a two-way table that shows the joint and marginal frequencies.

20. **HOW DO YOU SEE IT?** A research group surveys parents and coaches of high school students about whether competitive sports are important in school. The two-way table shows the results of the survey.

		Role		Total
		Parent	Coach	
Important	Yes	880	456	1336
	No	120	45	165
Total		1000	501	1501


- What does 120 represent?
- What does 1336 represent?
- What does 1501 represent?

21. **MAKING AN ARGUMENT** Your friend uses the table below to determine which workout routine is the best. Your friend decides that Routine B is the best option because it has the fewest tally marks in the “Does Not Reach Goal” column. Is your friend correct? Explain your reasoning.

	Reached Goal	Does Not Reach Goal
Routine A		
Routine B		
Routine C		

22. **MODELING WITH MATHEMATICS** A survey asks students whether they prefer math class or science class. Of the 150 male students surveyed, 62% prefer math class over science class. Of the female students surveyed, 74% prefer math. Construct a two-way table to show the number of students in each category if 350 students were surveyed.

23. **MULTIPLE REPRESENTATIONS** Use the Venn diagram to construct a two-way table. Then use your table to answer the questions.



- What is the probability that a randomly selected person does not own either pet?
 - What is the probability that a randomly selected person who owns a dog also owns a cat?
24. **WRITING** Compare two-way tables and Venn diagrams. Then describe the advantages and disadvantages of each.
25. **PROBLEM SOLVING** A company creates a new snack, N, and tests it against its current leader, L. The table shows the results.

	Prefer L	Prefer N
Current L Consumer	72	46
Not Current L Consumer	52	114

The company is deciding whether it should try to improve the snack before marketing it, and to whom the snack should be marketed. Use probability to explain the decisions the company should make when the total size of the snack’s market is expected to (a) change very little, and (b) expand very rapidly.

26. **THOUGHT PROVOKING** Bayes’ Theorem is given by

$$P(A|B) = \frac{P(B|A) \cdot P(A)}{P(B)}$$

Use a two-way table to write an example of Bayes’ Theorem.

Maintaining Mathematical Proficiency

Reviewing what you learned in previous grades and lessons

Draw a Venn diagram of the sets described. (*Skills Review Handbook*)

- Of the positive integers less than 15, set A consists of the factors of 15 and set B consists of all odd numbers.
- Of the positive integers less than 14, set A consists of all prime numbers and set B consists of all even numbers.
- Of the positive integers less than 24, set A consists of the multiples of 2 and set B consists of all the multiples of 3.