# 6.5 Properties of Logarithms

For use with Exploration 6.5

**Essential Question** How can you use properties of exponents to derive properties of logarithms?

Let  $x = \log_b m$  and  $y = \log_b n$ .

The corresponding exponential forms of these two equations are

 $b^x = m$  and  $b^y = n$ .

**EXPLORATION:** Product Property of Logarithms

Work with a partner. To derive the Product Property, multiply m and n to obtain

 $mn = b^x b^y = b^{x+y}.$ 

The corresponding logarithmic form of  $mn = b^{x+y}$  is  $\log_b mn = x + y$ . So,

 $\log_b mn =$  \_\_\_\_\_. Product Property of Logarithms

2

#### **EXPLORATION:** Quotient Property of Logarithms

Work with a partner. To derive the Quotient Property, divide *m* by *n* to obtain

$$\frac{m}{n}=\frac{b^x}{b^y}=b^{x-y}.$$

The corresponding logarithmic form of  $\frac{m}{n} = b^{x-y}$  is  $\log_b \frac{m}{n} = x - y$ . So,

 $\log_b \frac{m}{n} =$  \_\_\_\_\_. Quotient Property of Logarithms

3

#### **EXPLORATION:** Power Property of Logarithms

Work with a partner. To derive the Power Property, substitute  $b^x$  for *m* in the expression  $\log_b m^n$ , as follows.

$\log_b m^n = \log_b \left( \boldsymbol{b}^x \right)^n$	Substitute $b^x$ for $m$ .
$= \log_b b^{nx}$	Power of a Power Property of Exponents
= nx	Inverse Property of Logarithms

**174** Algebra 2 Student Journal

# 6.5 **Properties of Logarithms** (continued)

## **3 EXPLORATION:** Power Property of Logarithms (continued)

So, substituting  $\log_b m$  for x, you have

 $\log_b m^n =$  \_\_\_\_\_. Power Property of Logarithms

## Communicate Your Answer

- 4. How can you use properties of exponents to derive properties of logarithms?
- **5.** Use the properties of logarithms that you derived in Explorations 1–3 to evaluate each logarithmic expression.
  - **a.**  $\log_4 16^3$  **b.**  $\log_3 81^{-3}$
  - **c.**  $\ln e^2 + \ln e^5$  **d.**  $2 \ln e^6 - \ln e^5$
  - **e.**  $\log_5 75 \log_5 3$  **f.**  $\log_4 2 + \log_4 32$

## 6.5 Notetaking with Vocabulary For use after Lesson 6.5

In your own words, write the meaning of each vocabulary term.

base

properties of exponents

**Core Concepts** 

#### **Properties of Logarithms**

Let *b*, *m*, and *n* be positive real numbers with  $b \neq 1$ .

**Product Property**  $\log_b mn = \log_b m + \log_b n$ 

**Quotient Property**  $\log_b \frac{m}{n} = \log_b m - \log_b n$ 

**Power Property**  $\log_b m^n = n \log_b m$ 

Notes:

## Change-of-Base Formula

If a, b, and c are positive real numbers with  $b \neq 1$  and  $c \neq 1$ , then

$$\log_c a = \frac{\log_b a}{\log_b c}.$$

In particular,  $\log_c a = \frac{\log a}{\log c}$  and  $\log_c a = \frac{\ln a}{\ln c}$ .

#### Notes:

# 6.5 Notetaking with Vocabulary (continued)

## **Extra Practice**

In Exercises 1–4, use  $\log_2 5 \approx 2.322$  and  $\log_2 12 \approx 3.585$  to evaluate the logarithm.

**1.**  $\log_2 60$  **2.**  $\log_2 \frac{1}{144}$  **3.**  $\log_2 \frac{12}{25}$  **4.**  $\log_2 720$ 

In Exercises 5–8, expand the logarithmic expression.

**5.** 
$$\log 10x$$
 **6.**  $\ln 2x^6$  **7.**  $\log_3 \frac{x^4}{3y^3}$  **8.**  $\ln \sqrt[4]{3y^2}$ 

In Exercises 9–13, condense the logarithmic expression.

**9.**  $\log_2 3 + \log_2 8$  **10.**  $\log_5 4 - 2 \log_5 5$  **11.**  $3 \ln 6x + \ln 4y$ 

**12.** 
$$\log_2 625 - \log_2 125 + \frac{1}{3} \log_2 27$$
 **13.**  $-\log_6 6 - \log_6 2y + 2 \log_6 3x$ 

## 6.5 Notetaking with Vocabulary (continued)

In Exercises 14–17, use the change-of-base formula to evaluate the logarithm.

**14.**  $\log_3 17$  **15.**  $\log_9 294$  **16.**  $\log_7 \frac{4}{9}$  **17.**  $\log_6 \frac{1}{10}$ 

**18.** For a sound with intensity I (in watts per square meter), the loudness L(I) of the sound (in decibels) is given by the function  $L(I) = 10 \log \frac{I}{I_0}$ , where  $I_0$  is the intensity of a barely audible sound (about  $10^{-12}$  watts per square meter). The intensity of the sound of a certain children's television show is half the intensity of the adult show that is on before it. By how many decibels does the loudness decrease?

19. Hick's Law states that given n equally probable choices, such as choices on a menu, the average human's reaction time T (in seconds) required to choose from those choices is approximately T = a + b • log<sub>2</sub>(n + 1) where a and b are constants. If a = 4 and b = 1, how much longer would it take a customer to choose what to eat from a menu of 40 items than from a menu of 10 items?